

Diseases and Pests of Tea: Overview and Possibilities of Integrated Pest and Disease Management

Heinrich Lehmann-Danzinger*

Keywords: Tea, Pests, Diseases, Integrated Pest Management, Epidemiology, Root Diseases, Blister Blight, *Exobasidium vexans*, Scarlet Tea Mite, *Brevipalpus phoenicis*, Phytophagous Mites, Natural Enemies, Insecticides, Fungicides

Abstract

From the viewpoint of plant protection (phytomedicine), the demand for tea free from pesticide residues is in conflict with the demand for high quality from the consumer, and with the demand for high yield and low labour input, from the producer. High yield in tea has been mainly achieved by the elimination of tree shade, and reduction of losses due to diseases and pests with pesticides. Elimination of shade changed the agro-environment, with increased growth of weeds, higher input of fertilizers, and with increased susceptibility to certain diseases (stem canker due to sun-scorch, grey blight) and pests (mites). However, shade also can be detrimental to yield with unsuitable tree species, or too much shade in the rainy season which encourages the incidence of blister blight (*Exobasidium vexans*) and some insect pests (tea mosquito bug). Economic loss of tea due to diseases is higher compared to animal pests (pests), the blister blight being the main disease. Pressure of diseases and pests on tea depends also on the control strategy and the climatic environment. Reducing pesticides may be feasible by reducing the threat of diseases and/or pests through cultivation in less disease and pest prone environments (altitude, shade), in choosing disease or pest tolerant clonal tea-varieties, by choosing pesticides with low interference on natural enemies of pests and diseases, and by applying pesticides according to economic threshold. The long standing time of the tea bushes favours slow developing root diseases. These are reviewed and the integrated control is discussed in detail with eradication, with pesticides, and with improvement of soil microflora. Leaf and stem diseases are reviewed. Integrated pest and disease management of blister blight and the scarlet tea mite (*Brevipalpus phoenicis*), are discussed in detail.

* Dr. Heinrich Lehmann-Danzinger, Institut für Pflanzenpathologie und Pflanzenschutz, Georg-August Universität Göttingen, Grisebachstr. 8, D-37077 Göttingen, Germany. E-mail: lehman@gwdg.de

1 Introduction

Tea¹ originates in the mountains of Southeast-Asia. It is a small evergreen tree, which reaches a height of 10-15 m without cutting. In cultivation it is pruned down to 0.5 to 1.5 m in height to allow easier picking of the leaves. There are different tea varieties, but the most important are *Camellia sinensis* var. *sinensis* and *C. sinensis* var. *assamica*. The former has small, hard and serrated leaves, is more temperature tolerant (it can withstand -5°C), more aromatic, but has lower yields. The latter originates in south-east India (Assam), has large soft and smooth edged leaves, is mildly aromatic, high yielding and cannot withstand even the slightest frost. Nowadays, hybrids of the two varieties are mostly grown. A newer development is the selection of clones (see section "Blister blight of tea"), which are multiplied by cuttings.

Most of the world tea production² is produced in Asia (81.4%), second is Africa (15.5%), which is followed by Latin America (2.0%) (FAO, 1999). Tea is best grown in deep and well drained acidic soils (pH 4.5-5.0), well distributed rainfall of 1500-2000 mm per year, mild average temperatures of 18°-20°C and sufficient sunshine. These ecological conditions are prevalent in the tropics at higher altitudes between 800-2000 m (1400-2100 near the equator). According to Carr (1972) minimum rainfall for commercial yields is 1150-1400 mm/year, depending on how it is distributed. The temperature should not fall below 13°C and should not exceed an average of 30°C. In the subtropics tea has a dormancy period during winter and also in dry periods. The optimum sunlight intensity for tea is 50% to 80% from full radiation (EDEN 1976; PURSEGLOVE 1968). Suppression of weeds is essential. Tea is sometimes interplanted with a nurse crop or with green manure. Deep cultivation which would damage the surface roots should not be done. Shade was common, but is now controversial. The plucking interval depends on the length of time needed for production of new shoots (flushing) and, depending on altitude, varies from 70 to 90 days.

The biggest challenge for tea growers nowadays is to produce tea without pesticide residues. There is a growing demand for residue free tea, or at least low pesticide residue tea, in the highly industrialised countries and this tea fetches a good price. From the viewpoint of plant protection (phytomedicine) it should not be too difficult to grow tea without the aid of pesticides in some regions, if it was not for the phytomedical incompatible demand of the market for quality and the demand of the producer for high yield with low labour input. Reduced labour intensity has been achieved, starting in the seventies, by replacing manual weeding with herbicides, thereby introducing the risk of herbicide residues in the plucked tea. High yield has also been established by eliminating tree shade, which increased the weed problem, the need for synthetic fertilizers and increased the incidence of some pests and diseases. However the need for shade of tea is controversial (EDEN, 1976). The economic loss of tea due to diseases is generally

¹ Tea, *Camellia sinensis* (L.) O. Kunze, of the family Theaceae

higher compared to losses due to animal pests (pests). The pressure of diseases and pests on tea depends on the disease and pest control strategy of the plantation and also on the agro-ecological environment, which is different in specific tea growing regions and which varies at different altitudinal levels within the regions. Therefore one way of reducing pesticide application is to escape excessive pressure from specific diseases or pests by growing tea in environments less favourable to pests and diseases, which may not be always possible. Other alternatives are discussed below. Tea is a plantation culture, with a single tea bush remaining for a very long time at the same site (> 50 years). This might be desirable from an economic viewpoint, but not so from a phytomedical viewpoint. The healthy survival of the tea bush for a long time at the same site requires a good equilibrium (degree of integration) with the environment, which due to the management practices is not possible in highly efficient plantations.

Shade in tea through shade trees was common in the past. However, cultivating tea in full sunlight or with reduced shadow is more common now, since yield is higher and disease severity of the main leaf disease in Asia, the blister blight (*Exobasidium vexans*), is lower, this applies also to the attack of the tea mosquito bug (*Helopelthis* sp.). On the other hand, leaves falling from shade trees (acting as a nutrient pump) have a favourable impact on organic matter in the soil, which is important for the diversity and population density of the soil microflora and fauna. Spreading and incidence of root diseases and also nematodes is favoured by a disturbed soil microflora with a low species diversity and a low population density of antagonists to the fungal root disease agents.

The lack of equilibrium with the environment shows up in a number of diseases and pests, the decline diseases being an important disease group. The long standing time of the tea bushes is favouring slow developing root diseases spreading from one bush to another by root contact. In time, big dead patches or clearings due to these diseases appear in the plantations. The most important fungal diseases of tea, besides blister blight of leaves, are root diseases (Table 2).

2 Overview tea diseases

Since the leaves are the harvest product in tea, leaf diseases play an important role. The most important disease being the blister blight (Table 1) found in Asia, followed by the worldwide present root diseases (Table 2 and Section 2.1).

The information in Tables 1 and 2 has been recompiled from various sources, mainly Hainsworth, (1952); Eden, (1976); Semangun, (1983); Waller & Holderness (1997), and supplemented by the author's experience in Java, in Vietnam, in Peru, and Bolivia.

² World Tea production was 2 963 000 metric tons in 1998

Table 1: Leaf and stem diseases of tea

Common name	Causal agent & geographic distribution	Symptoms
blister blight (E) Blasenbrand (D) cacar teh (BI) quema de ampolla (S)	<i>Exobasidium vexans</i> Masee (Basidiomycotina, Ustomycetes, Exobasidiales) Asia only	First, pale yellow translucent spots, then circular blisters on leaf underside. Then white velvety and later circular brown spot
black blight, thread blight (E) Fadenbrand (G) mal de hilachas (S)	<i>Corticium koleroga</i> , <i>C. invisum</i> <i>syn. Pellicularia koleroga</i> (Basidiomycotina, Hymenomycetes, Aphyllophorales, Corticiaceae). World-wide	Leaves and twigs turn brown. Dead leaves are hanging on thin threads from the branches. Under humid conditions in the tropics
grey blight (E) Grauer Brand (D) quema gris (S)	<i>Pestalotiopsis theae</i> (Deuteromycotina, Coelomycetes) World-wide	Round, irregular, grey and necrotic leaf lesions. In the centre of older spots black fruiting bodies (acervuli). Mostly attacks weak or damaged tea bushes
red rust (E) Roter Rost (D) roya roja (S)	<i>Cephaleurus parasiticus</i> Karst (Algae, Heterokontophyta [=Chrysophyta]) World-wide	Red film covering leaves and branches
pink disease (E) Rosarote Zweigkrankheit (D) mal rosado (S)	<i>Corticium salmonicolor</i> Berk. & Broome (Basidiomycotina, Hymenomycetes, Aphyllophorales, Corticiaceae). World-wide	Pink crust on twig and branches. The portion above the branches dies off and cause dieback. Under humid conditions in the tropics
hypoxylon wood rot, wood rot (E); Holzfäule (D) pudrición de madera (S)	<i>Hypoxylon serpens</i> (Prs. ex Fr.) Ascomycotina, Pyrenomycetes) South India, Kenya	Stem canker. Wood with blackish patches and lines
low-country stem canker (E) Stamkrebs (D)	<i>Macrophoma theicola</i> (Petch) (Deuteromycotina, Coelomycetes) Sri Lanka	Stem-canker
thorny stem blight (E) Stacheliger Stammbrand (D) quema espinosa del tronco (S)	<i>Tunstallia aculeata</i> Petch (Ascomycotina, Sphaeriales). India	Branches die off. Fruiting bodies (perithecia) project from bark of dead branches giving a thorny appearance. Weak wound parasite

(E): English, (D): German, (BI): Bahasa Indonesia, (S): Spanish

Regarding leaf diseases, the black blight or thread blight (*Corticium invisum*), the pink disease (*Corticium salmonicolor*) and algae on leaves causing the "red rust" (*Cephaleurus parasiticus*), can be a severe problem in humid climates. The blister blight is discussed in detail in section 2.2.

Acute defoliation by the black blight is controlled by pruning the affected bushes, and burning the shoots on the spot. Plucking tea of the affected fields only in dry weather avoids disease transmission. New growth is protected with fungicide sprays (copper or organic fungicides, see section 2.2). *C. invisum* is also controlled by antagonists on leaf surfaces (phylloplane), such as the bacterium *Bacillus subtilis* (BARTHAKUR et al., 1993), which may provide a biological control agent for black rot. The grey blight can be controlled to a great extent with resistant varieties, such as IRB88 from Yakubita variety (Japan), CP-1 and TV-26 in SE-Asia, and 31/11 and 303/199 in Africa (Kenya). Red rust is controlled with repeated copper sprays and resistant clones (TV17, TV23, TV24, TV25, TV26, TV27) (BARTHAKUR et al. 1992).

2.1 Root diseases

Name and causal agents of root diseases: Several fungal species. See Table-2.

Economic importance: Hainsworth (1950) calculates the loss involved with root diseases, concluding that with a 1% loss of bushes, the loss of the potential crop expected in the tenth year would be 10%. In Africa, root diseases are the limiting factor of the tea production in many countries. In Kenya, losses up to 50% of tea bushes occurred (ONSANDO et al., 1997).

Symptoms

The general symptoms of root diseases is a dieback of twigs and branches of the bushes, and a wilt of the seedlings. In tea the first symptom of root diseases is wilting, which is not always clearly visible, and shows up first at noon. Then comes sudden yellowing or bronzing of leaves, followed by dieback (tip of branches begin to die), which can also show up as a sudden death of the whole bush. A characteristic of root diseases is the sudden appearance of symptoms. Most of the fungal dieback diseases (with exemption of the seedling diseases) have an incubation period of more than one year, meaning that it takes more than one year from infection to symptom expression.

Following the first symptom of dieback (wilting, bronzing, yellowing, or dying of tip of branches), the tea bush is killed in a relatively short time. This happens especially in the drier season, in which water must be supplied at a higher rate to the leaves by the root and vascular system. Leaves and twigs wilt first and die thereafter if the water supply rate is reduced, and water demand is high, due to low air humidity and high temperature. Reduced water supply may be either due to a limited uptake ability of the rotten roots, or to a reduced capacity of the vascular system, blocked by the dieback fungus.

Table 2 : Serious root diseases of tea

Common name	Causal agent	Symptoms on tea
red root rot (E) Rote Wurzelfäule (D) akar merah (BI) raíz roja (S)	<i>Ganoderma pseudoferreum</i> (Wakef.) van Overh. et Steinm; <i>G. philippi</i> (Basidiomycotina, Hymenomycetes, Aphyllophorales, Ganodermataceae)	Wilt of the plant and dieback. Roots with white surface mycelium which later turns red, to which soil adheres. Rot pale brown and hard, later pale buff and either dry or spongy
brick-red root rot (E) Ziegelrote Wurzelfäule (D) akar merah buta (BI) pudrición de rojo ladrillo (S)	<i>Poria hypolateritia</i> Berk. (Basidiomycotina, Aphyllophorales, Polyporaceae). Also <i>Ganoderma pseudoferreum</i> (see above),	Root surface white speckled with mycelial strands. These turn into a smooth sheet hardening into plates or ropes of red colour. They show up with scraping or washing
black root rot (E) Schwarze Wurzelfäule (D) akar hitam (BI) pudrición negra (S)	<i>Rosellinia arcuata</i> Petch; <i>R. bunodes</i> (Berk. et Broome) Sacc. (Ascomycotina, Pyrenomycetes, Sphaeriales, Xylariaceae)	Rot of root and stem base, black wood discolouration. White mycelia on roots, later grey to black. Ringing of stem. <i>R. bunodes</i> wide host range
charcoal stump rot, charcoal root, ustulina charcoal rot (E) Wurzelhalsfäule (D) leher akar (BI) pudrición de cuello (S)	<i>Ustulina deusta</i> (Hoffm.) Lind; <i>U. zonata</i> (Lév.) Sacc. (Ascomycotina, Pyrenomycetes, Sphaeriales). World-wide	No surface mycelium. Under bark of roots white fan-like patches. wood at base of stem with irregular double lines. Wide host range
diplodia root rot (E) Diplodia-Wurzelfäule (D) akar Diplodia (BI) pudrición radicular de diplodia (S)	<i>Lasiodiplodia theobromae</i> (Pat.) Griffon et Maubl. (syn. <i>Botryodiplodia theobromae</i> Pat.) (Deuteromycotina, Coelomycetes). Pantropical	Dieback. Root and stem-base rot. Blackened vascular-system and black discolouration of wood. Large number of host plants
armillaria root rot, root splitting disease (E) Hallimasch Wurzelfäule (D) pudrición de armillaria (S)	<i>Armillariella mellea</i> Vahl. Fr. (syn. <i>Armillaria mellea</i>) (Basidiomycotina, Hymenomycetes, Agaricales). Severe in Africa and Indonesia, in India rare	Sudden browning of leaves. Root splitting. White mycelial mat under the bark of stem base and roots
white root rot (E) Weiße Wurzelfäule (D) akar putih (BI) pudrición blanca (S)	<i>Rigidoporus lignosus</i> (Klotzsch) Imazeki (Basidiomycotina, Hymenomycetes, Aphyllophorales, Polyporaceae)	Dieback, white mycelium (rhizomorphs) on root. Roots rotten and white. Living wood decaying fungi

Table 2 (continued): Serious root diseases of tea

Common name	Causal agent	Symptoms on tea
brown root disease (E) Braune Wurzelfäule (D) pudrición radicular café (S)	<i>Phellinus noxius</i> (Corner) Cunn. (Syn. <i>Fomes noxius</i> Murr.) (Basidiomycotina, Aphyllophorales, Hymenochaetaceae). Africa, SE Asia, Australasia	Adherence of a crust of earth and gravel round the entire root. White or brown mycelium under the bark. Pale dry rot of wood
rhizoctonia seedling blight (E) Rhizoctonia Sämlingsbrand (D) quemadura de plántula por rhizoctonia (S)	<i>Rhizoctonia bataticola</i> (Taub.) Butl. (Deuteromycotina, Agonomycetes); teleomorph probably <i>Tanatephorus cucumeris</i> (Basidiomycotina, Hymenomycetes)	Seedlings wilt and dieback. Wide host range. Soil inhabiting fungi
seedling disease (E) Sämlingskrankheit (D) penyakit pesemaian (BI) enfermedad de plántula (S)	<i>Cylindrocladium ilicicola</i> (Deuteromycotina, Hyphomycetes), teleomorph <i>Calonectria</i> (Ascomycotina, Hypocreales)	Seedlings wilt and dieback. Pantropical, wide host range

(E): English, (G): German, (BI): Bahasa Indonesia, (S): Spanish,

Biology

Most of the root rotting fungi belong to the subdivision basidiomycotina and the rest to the ascomycotina, exempting *Lasiodiplodia theobromae* and *Fusarium oxysporum* (the latter not listed in Table 2), which belong to the deuteromycotina (fungi imperfecti), lacking characteristic visible fruiting bodies. The best way to identify the causal agents of the root and/or wood rot is by the fruiting bodies³. Unfortunately these often appear when the bush is already dead, or after incubation of samples with symptoms in a moist chamber (plastic bag). But, as described in Table 2, the individual fungal root rot species, can be recognised often by the symptoms on the roots and in the wood (Table 2). Identification based on pure cultures isolated from diseased tissue, is recommended and necessary with fungi belonging to the subdivision deuteromycotina. The fungal disease agents of root dieback have a wide host range on roots of wooden plants and are facultative soil pathogens, meaning that they can survive in soil without a host plant.

Epidemiology

In soil, other soil inhabiting micro-organisms keep the population of the facultative pathogens low. A higher diversity and population density of the saprophytic soil micro-organism will reduce the population of the facultative pathogens accordingly, hereby

³ The only reliable way of identifying fungal species is with taxonomy by fruiting bodies

lowering the chance of the facultative pathogens to attack healthy tea bushes. However, facultative root pathogens have very favourable growth conditions on root and wood residues in the soil, remaining from former trees and bushes of the site. These residues give rise to locally high populations of the facultative pathogens, meaning a high inoculum density. On such sites the facultative root pathogens can easily attack healthy roots of tea bushes.

Working on diagnostic and epidemiology of root rot on newly planted coffee, citrus and mango in Central America (Costa Rica, Nicaragua, El Salvador) at altitudes up to 800 m, the author found that dieback by black root rot (*Rosellinia* sp.) on coffee, and diplodia root rot (*Lasiodiplodia theobromae*) on citrus and mango, originated invariably through contact with buried residual wood or roots (mostly with visible mycelia). In older coffee plantations, infection occurred through the growth of mycelia or rhizomorphs from roots of infected neighbouring coffee bushes. Frequent downhill spread of the disease in up to 5-year-old coffee, may be due to inoculum (mycelia, spores?) spread by water (LEHMANN-DANZINGER, 1990, 1997). Dieback on rubber (*Hevea brasiliensis*), attacked by the black root rot (*Rosellinia* sp.) in South Kalimantan (Indonesia), was also due to root contact with wood debris. All these cases had, besides wood and root residues in the soil, other features in common: Soils with low pH, low content of organic matter, and previous clearing from more or less degraded forests. A disequilibrium with the environment and/or, in the case of *Hypoxylon serpens*, wounded stems, are also in tea responsible for attack by stem canker, caused by fungi such as *Macrophoma theicola* and *H. serpens*. Low organic matter leads to low diversity and density of soil microorganisms, meaning a disequilibrium of the soil environment.

A good example for a disequilibrium of the environment causing diseases, are the cork oaks: The Marmora forest in Morocco is composed of cork oaks (*Quercus suber*) as the dominant species. Within the past decades this forest has been overgrazed severely, damaging and stressing the oaks and hindering the renewal of the forest. The result was over-aging of the cork oak population. These oaks were severely attacked by the charcoal disease (*Hypoxylon mediterraneum*, Ascomycotina: Sphaeriales), causing stem canker and killing an appreciable number. The dead wood was then colonized by soil-borne saprophytic and facultative pathogenic wood decaying fungi: *Heterobasidion annosum*, *Armillariella mellea* and the propagules of these fungi accumulated in the soil. As a result newly planted acacias (*Acacia mollissima*) on part of the former forest, were attacked so heavily by root rot, that after 4 years over 50% of the trees had died (LEHMANN-DANZINGER, 1986). Similar high losses occurred in Kenya on tea attacked by Armillaria root rot, transmitted by buried wood or root residues (ONSANDO et al., 1997).

Control

From the biology and epidemiology of root diseases two preventive control strategies emerge: First, removal of stumps, other wood residues and large roots of woody plants from the soil prior to planting. This applies also to shadow trees removed from the

plantation. The second strategy is increasing the diversity and population density of the soil microflora, by augmenting the organic matter of the soil either with green manure, with appropriate shadow trees, or by adding organic manure such as compost. Woody plant residues should be destroyed by burning or burying them beyond reach of roots in the soil.

Controlling the root and dieback diseases appearing in the plantation is done by first removing the diseased bush and all neighbouring tea bushes together with their roots and destroying them by burning or burying them outside the plantation. Replanting is possible only after a two year fallow with non host plants, such as grass. If after two years there are still appreciable wood residues remaining in the soil, a one year treatment of the replanted tea bushes with systemic fungicides is recommended. Suitable systemic fungicides are the group of demethylation inhibitors (DMI)⁴, such as bitertanol, flusilazole, propiconazole, penconazole, hexaconazole and others, and in the group of morpholines tridemorph⁵. In Sri Lanka good results have been obtained by drenching systemic fungicides in 3-4 month intervals for a period of one year (ARULPRAGASAM, 1987). Applying fungicides of the same group consecutively should be avoided as to not promote resistance of the fungus against the fungicide. The two year waiting period can be shortened to one month, if soil fumigants are applied. The disadvantage of the available soil fumigants, such as chloropicrin (tear gas), the now banned methyl bromide, and formalin (formaldehyde), is that they are rather expensive (exempting formaldehyde), that covering of the site with a plastic sheet is necessary and that they have to be applied with the necessary precaution as not to harm the personnel involved. Formaldehyde is not as effective as methyl bromide. However it is less costly, easier to apply and a good fumigant when treated surfaces or products are wet. It acts in the gas phase and comes as a liquid with 36% or 40% formaldehyde. The dose for soil fumigation is 1.5 litres of a 40% product for each 10 m². Application is by diluting the formaldehyde in water and then spraying it with a can over the soil. Afterwards sealing with a plastic sheet for 77 hours is necessary (LEHMANN-DANZINGER, 1994). Following fumigation a waiting time of 3 weeks should be observed, before replanting. In any case application of organic manure (compost) is recommended before replanting.

2.2 Blister blight of tea

Name of the disease: Blister blight (E), cacar teh (BI), quema de ampolla (S), Blasenbrand (G).

Causal agent: *Exobasidium vexans*

⁴ The mode of action of the DMI's is inhibition of sterol biosynthesis

⁵ For an overview and an application oriented description of fungicides see LEHMANN-DANZINGER, 1994. A detailed description of fungicides is given by LYR, 1995

Geographic distribution: Indian subcontinent, south-east Asia and east Asia.

Economic importance: Losses due to blister blight extend up to 43% on annual crop basis. Heavy attack can result in death of the whole plant. Even with fungicide control 11.3% yield losses occur in Sri Lanka, during the peak attack at the altitudinal range of 1500m - <1800 m (FUCHS, 1989).

Symptoms: Symptoms appear first on young leaves and later on the older leaves, but only young leaves can be infected. First, pale yellow translucent spots appear, then circular blisters on the leaf underside with a dark green watersoaked zone around. Finally the blister becomes white and velvety (spore release). Following sporulation, the blister turns into a circular brown spot. Infected young stems are bent, distorted or girdled and may break off and die.

Biology: The causal agent *Exobasidium vexans* Masee belongs to the division Basidiomycotina, class Hymenomycetes, and here in the order Exobasidiales of the subclass Holobasidiomycetidae. The first symptoms appear following a 9-10 day incubation period. This period starts with the arrival of the spores on the leaf (inoculation). The first symptoms are irregular pale yellow translucent (watersoaked) spots which are clearly distinguishable on the underside of young leaves. Following 16 days after inoculation the spots become circular and the epidermis at the center becomes indented, forming a blister on the underside of the leaf. At 18-20 days after inoculation, the blister becomes first grey and then turns white and velvety. Here the lower epidermis of the blister is ruptured and spore bearing upright filaments (4-spored basidia) appear. Ripe spores are ejected with force from the basidia (GADD, 1949) and the lesion sporulates for about 1 week. Later on, the blisters turn brown and break up. The lesions appear then as circular spots with a dark brown centre surrounded by a light brown (tan) zone. The spores are thin walled, susceptible to desiccation, and direct sunlight. Therefore they can survive only a short time in dry air, and spread of the disease is difficult in the dry season. Surviving of *E. vexans* in the dry season is probably with immature lesions on leaves protected from direct sunshine.

Epidemiology: The blister blight was first reported from the Assam valley in 1868, at the border to Myanmar, where it caused little damage. Then it spread in 1908 to Darjeeling and from there in 1946 to south India and Sri Lanka. It also appeared in 1912 in Japan and 1930 in Vietnam. In 1949 the blister blight jumped over to Sumatra and from there in 1950 to neighbouring Java (Indonesia). Now it is endemic throughout most tea-growing areas of Asia, where it has reached epidemic proportions. It has not been reported in Africa or America. How spreading occurred in India and Asia still remains an unresolved problem. On opinion of the author this must have happened by leaves with immature lesions, probably with transport of tea seedlings.

Depending on climatic conditions, the full reproduction cycle (from inoculation and infection to release of spores) of *E. vexans* varies from 16 to 25 days. The disease

occurs mainly in the rainy season. Cool, moist, still air favours infections, as do higher elevations. According to FUCHS (1989), more than 4-5 hours of sunshine/day keeps attack below danger level. In Sri Lanka severe attack of blister blight occurs mainly above 600m of altitude, with the most heaviest attack at 1500m - <1800 m. A similar sharply increasing incidence of the yellow Sigatoka⁶, a banana and plantain leaf disease, occurred in the Andes mountain range⁷ at >800-<1800 m, with a peak at 1100m (LEHMANN-DANZINGER, 1987). In this altitudinal range condensing air leads to frequent fog, high humidity, and drizzling rain. These conditions lead to a dramatic decrease of sunshine and altogether favour fungal leaf diseases and, at the same time, lower the disease resistance in crops not adapted to the environment. This being the case with tea and bananas/plantains. Tea cultivated in the above range of altitude in south America is affected by losses due to diseases, such as algae, grey blight and pink disease. However, if blister blight is introduced there, losses are going to be very heavy.

SUGHIA (1997) found that blister blight was enhanced by the minimum and average temperatures and the relative humidity, but was negatively influenced by the average rainfall or hours of sunshine. Survival of the fungus in the off season was on necrotic blisters. The disease appeared then (end of May in Himachal Pradesh, India) on volunteer tea seedlings under mother bushes, and here the inoculum for infection in the following wet season was certainly build up. On opinion of the author of the present paper, *E. vexans* survives the dry season with slow developing or dormant immature lesions, which can develop further and sporulate when favourable environmental conditions appear.

Blister blight is more serious in plants recovering from pruning, since they are debilitated and have mainly young leaves. This susceptible period may extend up to 2 years after pruning. Pruning in the dry season avoids attack of the extremely susceptible leaves in the first period of recovery. Of course dry-weather pruning delays recovery and may cause sun-scorch of the stems (frame) and make them susceptible to stem cankers (caused by *Hypoxylon* and *Macrophoma*) and insect pests. The remedy to this drawback is providing shade during the dry season, which may be reduced (by pruning) afterwards in the rainy season, in order to reduce blister blight incidence. An alternative is pruning tea in the wet season and protect the bush from blister blight with systemic fungicides (see control).

Control: The most important factor contributing to the incidence and severity of blister blight is, besides temperature, humidity. Therefore the main attack on tea is in the rainy season. Newer tea clones, such as GMB-1, GMB-2 and GMB-3 show some resistance to the disease in comparison to older clones (TRI 2025). Clone PS1 is highly resistant in Indonesia (MARTSUPONO, 1991). The resistance transduces to a lower control level with

⁶The causal agent of the yellow Sigatoka disease is *Mycosphaerella fijiensis* (Ascomycotina)

⁷ The survey was conducted in Venezuela, Colombia, Peru and Bolivia, from which the latter two had tea plantations

fungicides, or even with no fungicide control. Therefore more attention should be paid to resistance against blister blight when planting tea at higher altitudes and breeding new clones.

Light pruning only in dry weather (addressed above) and reduction or removal of shade assist control with fungicides. The main strategy of control is prevention from infection. A second strategy is to start the wet season with an inoculum as low as possible and suppress sporulation of the fungus. The tea leaves are traditionally protected from infection of *E. vexans* by application of copper-oxichloride at 125g a.i./ha, or at higher concentration after pruning. Since only the young leaves and tender twigs can be infected by the fungus, only these have to be sprayed with the fungicide. The layer of copper on the leaves will kill the germinating spore of *E. vexans*. However, this protection is sometimes not enough during the rainy season. Here, cupric oxides give better results. Nickel chloride is superior to copper fungicides, since it has a curative action. To control the blister blight in India, a mixture of copper fungicide and nickel chloride is sprayed at 7-10 day intervals throughout the rainy season. However, copper fungicides stimulate the increase of the population of tea mites, especially the scarlet tea mite (Tenuipalpidae), which is the main pest of tea grown at altitudes below 1400 m, and also the purple mite and pink mite, belonging to the family Eriophyidae (Table 3).

An alternative to copper fungicides are the well established protective organic fungicides which are Calixin® (a. i. tridemorph)⁸, and Sicarol® (a. i. pyracarbolid)⁹. These are applied at considerably lower doses at 14 day intervals and do not seem to enhance the multiplication of mites. A disadvantage of piracarbolid is that it enhances the population of the mosquito bug (*Helopeltis* spp.) (Table 3). A curative effect is reported for tridemorph, meaning that it can stop further development of non sporulating lesions to sporulating lesions (see "biology"), another advantage is that it does not stimulate mites. A disadvantage, at least with Calixin®, is that it can be rapidly washed off by heavy rainfall during the monsoon. Therefore this fungicide should be applied at 14 day intervals during the beginning of the rainy season (2-3 applications), to lower the spore inoculum, and the end of the rainy season (3-4 applications), to keep the mite population down before the starting dry season (VENKATA RAM, 1980). A curative effect is achieved by the newer (and more expensive) group of organic fungicides with systemic action in the plant, and belonging to the group of the benzimidazole fungicides, especially Benlate® (a.i. benomyl), and to the demethylation inhibitors (DMI), especially Bayleton® (a.i. triadimefon), Tilt® (a.i. propiconazole) and Anvil® (a.i. hexaconazole) which can be applied up to 21 day intervals. These can also be mixed with copper oxychloride. Being systemic, these fungicides cannot be washed off by heavy rain, providing that there is enough time for penetration before rainfall after application. The curative treatment is important to reduce the spore inoculum by suppressing sporulation

⁸ a. i. = active ingredient of the commercial product

⁹ Pyracarbolid is no longer manufactured or marketed according to Tomlin (ed.), 1997

(by stopping lesion development), when control is not possible with the available protective fungicides. This is the case when the inoculum originates from heavily infested disease nests ("hot spots") in the plantation, these should then be treated selectively with curative fungicides. Selectively treating only the hot spots is an alternative to treatment of the entire tea plantation, if disease incidence is not too high (for instance at lower altitudes). This strategy would have less impact on beneficial entomopathogenic fungi in comparison to a broad application of copper or other fungicides in the plantation. Generally, the copper fungicides require application at rather high doses in comparison to organic non systemic fungicides and organic systemic fungicides. Applying copper fungicides frequently results in accumulation of copper in the soil which, at higher concentrations, becomes phytotoxic, especially in soils with a low content of organic matter.

Setting up a disease forecast model for blister blight should be possible by relating bioclimate to disease development. Bioclimate includes continuous measurements (at the height of the tea bushes) of temperature, relative humidity, leaf wetness, sunshine and rainfall. Disease development includes lesion density, spore density in the air, and follow-up of lesion development. Encouraging results have been achieved by Sugha (1997) (see epidemiology). Having set up the forecast model, blister blight severity forecast should be possible by measuring only the relevant parameters.

Conclusion: Controlling blister blight at higher altitudes in climates with a strong rainy season is not possible without fungicides. However, the number of applications and the amount of the applied fungicide can be lowered, as shown above. A problem is that pests are encouraged, mainly mites, by the sprayed fungicides. Enhancement of mites can be reduced or abolished by the selections of non interfering fungicides, and application of cuprous fungicides only in the middle of the rainy season, when mites are not active. The outlined integrated control strategy could further reduce or even abolish fungicide residues in tea.

3 Overview of tea pests

An overview of the main animal pests (pests) of tea is given in table 3 and table 4. Natural enemies in Asia are listed in Table 4. The information in Tables 3 and 4 has been recompiled from various sources, mainly Hainsworth, E. (1952); Krishna & Weesner (1970); Eden (1976); Das (1974); Sarma (1979); Kalshoven & Van der Laan (1981); Dharmadi (1988); and Muralcedharan (1992), and supplemented by the author's experience in Java (Indonesia), Vietnam and Peru (South America).

From the pests in the overview, phytophagous mites are considered one of the most limiting factors in tea production. The second most damaging pests are the *Helopeltis* bugs, which are also of economic importance in Africa. The third most important group of pests, listed in table 4, are leaf feeding caterpillars of the lepidoptera order, here

Table 3: Important pests of tea

Common name	Scientific name, order and family and geographic distribution	Symptoms, part of plant attacked
mosquito bug	<i>Helopeltis schoutedeni</i> Reut., E-Africa; <i>H. orophila</i> Ghesq., W-Africa. (Heteroptera: Miridae)	Brown centred black lesions followed by holes in the leaves. Scorching of leaves with heavy attack.
mosquito bug, tea bug	<i>Heliopeltis antonii</i> Signoret (Heteroptera: Miridae). Sri Lanka, India, Indonesia, Vietnam. Occurs from 200 to 1400 m altitude	Some species cause stem canker with black lesions on stems.
tea mosquito bug, cocoa mirid	<i>Helopeltis theivora</i> Waterhouse (Heteroptera: Miridae), Sri Lanka, India, Bangladesh, Indonesia). Occurs up to 600 m	The mosquito bug (probably <i>Helopeltis</i> sp.) is also present in Peru (South America)
scarlet mite, red-and-black-flat mite, red-crevice mite	<i>Brevipalpus phoenicis</i> Geijskes (Acarina: Tenuipalpidae) World-wide (tropics)	Attack of tea flushes and leaves, corky area on leaf undersides, leaves dry up. Attack below 1400 m of altitude.
carinate tea mite, purple tea mite	<i>Calcarus carinatus</i> Green, (Acarina: Eriophyidae). Sri Lanka, India, Indonesia, Malaysia, China	Distortion of young leaves
yellow tea mite	<i>Polyphagotarsonemus latus</i> (syn. <i>Polytarsonemus latus</i>) (Acarina: Tetranychidae, Tarsonemidae) World-wide	Flush leaves are cupped or otherwise distorted, corky brown area between main veins on leaf underside. Browning of flushes
red coffee mite	<i>Oligonychus coffeae</i> Nietner (Acarina: Tetranychidae) practically world-wide, tropics	Yellowish-brown, rusty or purple discolouration of leaves. Fully developed leaves are mostly attacked
pink tea rust mite	<i>Acaphylla theae</i> (Watt) (Acarina: Eriophyidae) Bangladesh, India, Malaysia, Indonesia, China, former USSR	
red coffee borer	<i>Zeuzera coffeae</i> Nietner (Lepidoptera: Cossidae). India, Sri Lanka, Malaysia, Indonesia	Larva red to violet-brown, often with yellowish rings. Larvae bore inside stems and branches. Dieback

Table 3 (Continued): Important pests of tea

Common name	Scientific name, order and family and geographic distribution	Symptoms, part of plant attacked
live wood termites	<i>Glyptotermes dilatatus</i> Bug. & Pop., <i>Postelectrotermes militaris</i> (Des) (Isoptera: Kalotermitidae) Sri Lanka	Attack of stems. Dieback. Colonies in the trunc region. Breeding and colony maintenance is more successful by <i>P. militaris</i> in tea stems
	<i>Microcerotermes</i> spp., (Isoptera: Termitidae). Fungivorous termites. India, Malaysia, Java	
ground termites	<i>Macrotermes gilvus</i> (Hag.) (Isoptera: Termitidae). Indonesia, Malaysia, Vietnam, Philippines	Nest in the ground. Attack of tea seedlings
tea shoot hole borer (stem borer)	<i>Xyleborus fornicatus</i> (Eichoff) India, Sri Lanka, SE Asia. <i>Xyleborus morigenus</i> (Blanford), Indonesia. (Coleoptera: Scolytidae)	Attack of stems. Dieback. Shoot holes in stem

Table 4: Serious insect pests of tea and natural enemies in Asia

Common name	Scientific name geographic distribution	Natural enemies present on tea & remarks
tea tortrix	<i>Homona coffearia</i> Nietner <i>Tortrix dinota</i> (in Malawi) (Lep.: Tortricidae) India (subcontinent), China	Larval parasitoid <i>Macrocentrus homonae</i> (Hym.: Braconidae). Introduced from Java successfully in Sri Lanka, controlling there <i>H. coffearia</i>
tea flushworm	<i>Cydia leucostoma</i> Meyrick (<i>Laspeyresia leucostoma</i>) (Lep.: Tortricidae) India, Bangladesh, Indonesia	Ectoparasite <i>Asymplesiella india</i> , <i>Elasmus homonae</i> (Hym.: Eulophidae); <i>Apanteles</i> sp. (Hym.: Braconidae)
tea leaf roller	<i>Caloptilia theivora</i> Walsingham (<i>Gracillaria theivora</i>) (Lep.: Gracillariidae)	<i>Asympleia india</i> (Hym.: Eulophidae) <i>Bethylus distingma</i> (Hym.: Bethyilidae)
tea leaf miner	<i>Tropicomyia theae</i> (Cotes) (Diptera: Agromyzidae)	India and Sri Lanka
bunch caterpillar	<i>Andraca bipunctata</i> Wlk. (Lep.: Bombycidae)	Parasitoid tachinid fly on 1st and 2nd larval instar. <i>Bacillus</i> sp. on late instars
tea looper	<i>Buzura suppressaria</i> (Guen.) (syn. <i>Biston suppressaria</i>). (Lep.: Geometridae). From Sri Lanka to China	Indian subcontinent, Indonesia, China

Table 4 (Continued): Serious insect pests of tea and natural enemies in Asia

Common name	Scientific name geographic distribution	Natural enemies present on tea & remarks
lobster caterpillar	<i>Staurops alternus</i> Walker (Lep.: Notodontidae)	<i>Apanteles taprobanae</i> (Hym.: Braconidae), <i>Carcelia guava</i> , <i>Zenillia sumatrensis</i> , <i>Z. casumatarensis</i> , <i>Tricholya sorbillens</i> (Dip.: Tachinidae)
nettle grubs	<i>Darna (Macrolepta) nararia</i> (fringed nettle grub), <i>Thosea sinensis</i> , <i>T. cervina</i> , <i>Setora nitens</i> , <i>Parasa</i> sp. (Lep.: Limacodidae)	<i>Bacillus thuringiensis</i> , granulosis virus, <i>Fornicia ceylonica</i> (Hym.: Braconidae), <i>Neoplectrus maculatus</i> , <i>Platyplectrus natadae</i> , <i>Metaplectrus thosaeae</i> , <i>Euplectrus</i> sp. (Hym.: Eulophidae); <i>Trichoglya sor-billans</i> (Dip.:Tachinidae)
jelly grub	<i>Cheromettia apicata</i> (Moore) (Lep.: Limacodidae)	<i>Batotheca nigriceps</i> (Hym.: Braconidae)
hemipteran pests	<i>Helopeltis</i> spp. (Heteroptera: Miridae) <i>Empoasca flavescens</i> (Auchenorrhyncha: Cicadellidae)	<i>Sycanus collaris</i> (Het.: Reduviidae), <i>Hierodula</i> sp. and <i>Tenodera</i> sp. (Man.: Mantidae), Wasp (Hym.: Drynidae)
tea aphid	<i>Toxoptera aurantii</i> Boyer (Hemiptera: Sternorrhyncha)	<i>Aphilius</i> sp. (Hym.: Aphelenidae) <i>Trioxys</i> sp. (Hym.: Braconidae) <i>Syrphus balteatus</i> (Dip.: Syrphidae)
tea shot-hole borer	<i>Xyborus fornicatus</i> (Eichoff) (Coleoptera: Scolytidae)	<i>Atherigona orientalis</i> (Dip.: Muscidae)
red coffee borer	<i>Zenzera coffeae</i> Nietner (Lep.: Cossidae)	<i>Bracon zeuzerae</i> , <i>Microbracon</i> sp. (Hym.: Braconidae), <i>Isoturnia chatterjeeana</i> , <i>Carcelia kockiana</i> (Dip.: Tachinidae)

All listed pests, except the tea shot-hole borer and the red coffee borer, feed on tea leaves

Lep. = Lepidoptera (moths, butterflies); Hym. = Hymenoptera (wasps); Dip. = Diptera (flies); Man. = Mantodea (praying mantids) (Blattopteriformia)

Geometridae and Limacodidae feed mainly on maintenance leaves. In Africa the mosquito bug *Helopeltis schoutedeni* is capable of causing annual losses up to 55% (RATTAN, 1992). The main pest of tea in Java and other regions of south-east Asia, from 400-1400 m altitude, is *H. antonii*. The adult bug is only 5 mm long and at a first view, resembles a mosquito¹⁰, however it is a bug having 2 pairs of wings and belongs to the

¹⁰ A true mosquito has only one pair of wings and is classified therefore in the order Diptera (flies)

order Heteroptera. In the frame of integrated control a wide range of measures have been applied in Java to control the mosquito bug (DHARMADI 1990a, 1990b). These measures include shortening of the plucking cycle, reducing nitrogen application, pruning in the correct time, control of broadleaved weeds, mechanical elimination of bugs with plucking, and use of released mantid predators (table 4). Application of insecticides becomes necessary when the above techniques are not sufficient to keep the pest below the economic threshold level of 8 adult *Helopeltis* bugs/2 m². This level reduces 78.75% of the leaf production within 8 weeks.

Mosquito bugs have been controlled with the organo chlorine insecticide gamma-HCH (γ -HCH) (= lindane). Nowadays the pyrethroid insecticide cypermethrin and permethrin (Ambush[®], Torpedo[®]), organo-phosphorous insecticides fenitrothion (Sumithion[®], Folithion[®]), and formothion (Anthio[®]) are preferred. However, formothion stimulates the scarlet mite (DHARMADI, 1988). A study by Deka et al. (1998) found that plant extracts of pongam (*Pongamia pinnata*) and wild sage (*Lantana camara*) have anti-feedant and repellent effects on the tea mosquito bug *H. theivora*. Tea shoots sprayed with a 5% chloroform extract of *P. pinnata*, showed only 43 spots on tea leaves caused by tea mosquito bug, in comparison to the untreated control with 653 spots 72 hours after treatment. This finding shows that an integrated control of mosquito bugs may be feasible by replacing insecticides with plant extracts. The further description and discussion of the mosquito bug might be desirable here, but must be postponed in favour of an in depth treatment of the mite problem in tea.

The natural enemies of the lepidopteran pests in tea are parasitoids¹¹ belonging either to the order hymenoptera (wasps), or diptera (flies) (table 4). This opens the possibility for control of the lepidopteran pests selectively with insecticides affecting only lepidoptera, such as the *Bacillus thuringiensis* (B.t.) based insecticides. B.t. insecticides have the advantage of not leaving residues on tea, if applied properly.

From the phytophagous mites, the scarlet mite is considered one of the most damaging and most difficult to control. It is dealt with in a separate section. Mites are active during dry weather. In south India 4 rounds of spraying acaricides were the minimum for mite control. Severe infestations require even 6 applications (SARMA, 1979). The acaricides used on tea worldwide are usually the organo phosphorous insecticide-acaricide dimethoate, and the organo chlorine acaricide-insecticide dicofol (see also Table 7). Both are broad spectrum acaricides-insecticides, meaning that they kill other mites and also insects besides the target mite. Therefore most of the natural enemies of mites (Table 5) and the other insect pests (Table 4) of the tea are killed by broad spectrum insecticides/acaricides, which promotes damage of tea by the other insect pests which are normally controlled by natural enemies (Table 4). Sarma (1979) reported that preda-

¹¹ Parasitoid is an animal (insect) that feeds and lives for a long time in or on another living animal (host insect), consuming all or most of its tissues and eventually killing it.

tor populations tended to decline after only one spray of the organochlorine insecticide chlorobenzilate¹², and nearly disappeared after several sprays.

Phytophagous mites have a number of natural enemies listed in Table 5, which under “normal” conditions (without the use of pesticides) control these mites on tea. The fungi

Table 5: Natural enemies of phytophagous mites on tea in India

Phytophagous mite species & family	Natural enemies	Order and family	Action of the natural enemy
red coffee mite	<i>Menochilus sexmaculata</i> (F.)	Coleoptera: Coccinellidae	predators
Oligonychus coffeae (Tetranychidae)	<i>Scymnus</i> sp.	(ladybird beetles)	
	<i>Verania vinta</i>		egg predators
	<i>Stethorus gilvifrons</i>		
	<i>Jauravia</i> sp.		
	<i>Oligota</i> sp.	Coleoptera: Staphylinidae (rove beetles)	predators
	<i>Orius</i> sp.	Heteroptera: Anthocoridae (minute pirate bugs)	
	<i>Micromus timidus</i>	Neuroptera: Hemerobiidae (brown lacewings)	
	<i>Phytoseiulus persimilis</i>	Acarina: Phytoseiidae	
carinate tea mite <i>Calcarus carinatus</i> (Eriophyidae)	<i>Typhlodromus (Amblyseius) ovalis</i>	Acarina: Phytoseiidae	
	other predatory mites	Acarina: different families	
scarlet mite <i>Brevipalpus phoenicis</i> (Tenuipalpidae)	predatory mites from different families	Acarina: Phytoseiidae, Trombiculidae, Anystidae	
	<i>Orius</i> sp.	Heteroptera: Anthocoridae (minute pirate bugs)	
	<i>Semidalis fletcheri</i>	Neuroptera: Coniopterygidae (dusty-wings)	
all mites	<i>Hirsutella</i> ?, <i>Penicillium</i> sp., <i>Aspergillus</i> sp., <i>Cladosporium</i> sp., <i>Botryosporium</i>	fungi	disease (entomopathogens)

Data mainly from Sarma (1979)

¹² The insecticide chlorobenzilate has been removed from the marketplace in the USA in 1989 and the registration was suspended in Germany. Now chlorobenzilate has been replaced in tea by the organo phosphorous insecticide trichlorphorhon (Dipterex®)

acting as entomopathogens (Table 5) are, with the exception of *Hirsutella*, not specific entomopathogens, but have been repeatedly isolated from dead phytophagous mites on tea. These and other fungi certainly play an important role in reducing the mite population in humid weather. The numerous predacious natural enemies of mites listed in Table 5 and 8 show that these, together with the listed pathogenic fungi, should be able to keep the mite population below the economic threshold level. It seems that this was the case before indiscriminate use of insecticides in tea because, before this time and according to a number of authors, phytophagous mites have never been a serious problem on tea. Of course, fungi are controlled by fungicides. So when unspecific broad action fungicides, such as copper based products, are applied to control the blister blight disease, the fungi acting as natural enemies are also killed. This effect has been well documented in south India (SARMA, 1979), where the use of residual copper sprays has increased the incidence of the red spider mite (*Oligonychus coffeae*) on tea, probably by killing the naturally occurring entomopathogenous fungi (described in Table 5).

The termite *Glyptotermes dilatatus* Bugnion & Popoff (Table 3) is a pest of tea in Sri-Lanka at middle altitudes - below 600 m (2000 feet). It has a broad host range attacking also coffee, rubber and cocoa. The soldiers are rather small of 7 mm length and the first stage larva has the sides of the thorax extended into 'wing-like' pads. The termite excavates longitudinal galleries and form a network of communicating tunnels, giving the wood a honey comb appearance in a longitudinal section of the nest in the stem. Roots are rarely attacked (KRISHNA & WEESNER, 1970). The termite *Postelectrotermes militaris* (Desneux) (*Neotermes militaris*) is a pest of tea in Sri-Lanka at higher altitudes occurring chiefly at 1200 to 1600 m. It destroys the tea bush, hollowing out the stem, branches, roots and even rootlets. The winged termite stage gains entry into the plant through dead snags and wood rot cavities, which should apply also to *Glyptotermes dilatatus*. Nesting occurs in the heartwood. The nest has no definite system of tunnels and chambers and the cavities appear as filled with earth. The final winged stage takes 7 years to develop and a single colony may inhabit a tea bush for over 15 years (BEESON, 1941).

3.1 Scarlet tea mite (*Brevipalpus phoenicis*)

Common name: Scarlet mite, red-and-black-flat mite, red-crevice mite.

Scientific name: *Brevipalpus phoenicis* (Acarina: Tetranychioidea, Tenuipalpidae).

Geographic distribution: Pantropical in most tea producing countries.

Host plants: Citrus (*Citrus* spp.), transmission of citrus leprosis disease; tea (*Thea*); coffee (*Coffea*) transmission of ringspot disease; date palm (*Phoenix dactilifera*); rose mallow (*Hibiscus rosa-sinensis*); passion fruit (*Passiflora edulis*) and 63 other host plant genera (PITCHARD & BAKER, 1958).

Economic importance: The scarlet tea mite is one of the main pests of tea in south-east

Asia during the six-month long dry season. It constitutes a serious pest problem in most tea producing countries (India, Sri Lanka, Bangladesh, Indonesia, Kenya, Malawi and Mauritius), where it often causes loss of 13% of the yield in tea. The injury and degree of infestation decreases with increasing altitude of the tea plantation, and is much lower above 1500 m of altitude. It seems that the mite problems are in some way related to the extensive use of pesticides. Copper fungicides, which are applied to tea to control the blister blight, stimulate the scarlet mite in tea (OOMEN, 1982). Also insecticides, which are applied to control the tea bug (*H. antonii*), stimulate the scarlet mite, especially the insecticide Anthio® (a. i. formothion). This is certainly due to the elimination of the mite's natural enemies. Herbicides decrease the population of the mite, most probably by killing secondary host plants of the scarlet mite (DHARMADI, 1988).

Symptoms and injury

The scarlet mite attacks the older leaves (tea maintenance leaves), their petioles and the non-lignified twigs throughout the year. The mites prefer to stay on the main veins of the leaf underside and crevices, especially on the leaf base and petiole. These parts become covered with a necrotic shade. With heavy attacks there is bronzing on the leaf underside, then the leaves fall off. Following damage to the leaves, the mites migrate upward to younger leaves. The attack is heavier on old tea bushes (>50 years) and debilitated plants. Therefore, some sections of a tea plantation might show a more severe attack than others.

The yellow tea mite *Polytarsonemus latus* attacks tea flushes in the rainy season. The flushes get brown and may die off. The economic losses are similar to those of the scarlet mite.

Table 6: Influence of the temperature on the duration of the life cycle of the scarlet mite *Brevipalpus phoenicis*

Host plant	Temp.	Egg [days]	Larva + protochrysalis [days]	Protonymph + leutochrysalis [days]	Deutonymph + teleichrysalis [days]	Total time 'egg-adult' [days]
tea (1)	19-23°C	14.4	5.4	6.3	7.4	33.5
tea	20-33°C	9.4	5.6	5.8	6.9	27.7
papaya	20°C	21.6	10.4	8.4	7.3	47.7
papaya	25°C	9.4	6.5	6.5	6.9	29.3

Adapted from Oomen (1982)

Table 7: Acaricides applied to tea to control mites (based on restrictions of tea importing countries)

Active ingredient common name	Activity	Commercial product (trade name)	Chemical group	Toxicity class EPA(4)	Maximum residue limit in tea
binapacryl	acaricide/ fungicide	Morncide®, Acracid®, Endosan®	Dinitro- phenol	II	0.3 mg/kg(3)
dicofol	acaricide	Kelthane®, Acarin®	Organochlo- -rine	II or III	2.0 mg/kg(2)
dimethoate	insecticide/ acaricide	Cygon®, Champ®, Perfekthion®, Rogor®, Roxion®, Tara	Organo- phosphate	II	0.2 mg/kg(2)
Chinomethionat oxythioquinox quinomethionat	fungicide/ acaricide	Morestan®	Organosul- phur	III	0.02 mg/kg(3)
cyhexatin (1)	acaricide	Plictran®, Mitacid®	organotin	III	2.0 mg/kg(2)

⁽¹⁾ cyhexatin (Plictran®) has been removed from the marketplace in the USA as a consequence of FIFRA Amendment's Reregistration Data Requirements

⁽²⁾ maximum residue limit for tea in Germany (PERKOW & PLOSS, 1999)

⁽³⁾ maximum residue limit in Germany, not extra specified for tea (PERKOW & PLOSS, 1999)

⁽⁴⁾ toxicity class equivalent to EPA (= Environmental Protection Agency, USA)

Biology

The scarlet mite is polyphagous and has many host plants such as the citrus and papaya (see section "host plants"). The adult of the scarlet mite measures only 0.28 x 0.16 mm and is therefore difficult to see with the naked eye. Adults eggs and larvae are red in colour.

They feed and maintain themselves on the leaf underside. The mite has five stages which are listed in Table 6. The egg and chrysalis stages are immobile and remain adhered to the leaf. The reproduction of the scarlet mite is parthogenetic¹³. Table 6 indicates that the duration of the life cycle of the scarlet mite is shortened by a higher ambient temperature. This means that multiplication is faster at lower altitudes, compared to higher altitudes. This explains the much lower infestation above 1500m of altitude. The population usually increases at the end of the dry season. The average multiplication rate (rate of increase) is 0.061, which is far less than for other spider mites.

¹³ Parthogenesis: Reproduction without sexual process (fertilization)

Table 8: Predators which prey upon the scarlet mite *Brevipalpus phoenicis* in Indonesia

Taxonomic group	Scientific name
Acarina: Phytoseiidae (predatory mites)	<i>Amblyseius deleoni</i> Muma et Denmark
	<i>Amblyseius largoensis</i>
	<i>Amblyseius tamatavensis</i> Blommers
	<i>Amblyseius</i> w (near <i>A. newsami</i>)
	<i>Amblyseius</i> x (near <i>Iphiseius</i>)
	<i>Amblyseius</i> z (<i>A. caudatus</i>)
Acarina: Stigmaeidae (predatory mites)	<i>Zetzellia</i> spp.
	<i>Zetzellia maori</i> Gonzalez
	<i>Agistemus</i> spp.
	<i>Agistemus terminalis</i> Quayle
	<i>Agistemus denotatus</i> Gonzalez
	<i>Agistemus feschneri</i> Summers
	<i>Agistemus arcypaurus</i> Gonzalez
Acarina: Podocinidae (predatory mites)	<i>Lasioseius</i> spp.
Diptera: Cecidomyiidae (gall midges)	<i>Lestodiplosis oomeni</i> Harris
Hemiptera: Anthocoridae (minute pirate bugs)	<i>Wollastionella testudo</i> Carayon

(based on data of OOMEN, 1982, HARRIS, 1982,)

Control

The economic threshold level of the scarlet mite is 25 eggs or adults per leaf, which cause a 13% yield loss (OOMEN, 1982). To control the scarlet mite most of the tea plantations have to be treated up to three times per year with the acaricides listed in Table 7. A control based on integrated pest management (IPM) is based first on assessments of the mite damage or of the mite population (if visible) during plucking (harvest) and then on the treatment with acaricides of the hot spots of the plantation, in which the damage or insect population is high. The acaricides listed in Table 7 are available for control of the scarlet mite on tea. Surprisingly the pyrethroid insecticide acaricide permethrin¹⁴ and the fungicides mancozeb and maneb are also toxic to the scarlet mite, reducing significantly their survival, oviposition and multiplication. The mite itself is not affected by copper fungicides (copper oxy-chloride, copper oxyde) and the insecticide carbaryl, belonging to the group of carbamates. As stated before, copper fungicides stimulate mites, the cause is not known exactly, but possible explanations are that cop-

¹⁴ Maximum residue limit in tea is 0.05mg/kg in Germany

per fungicides kill predators, they stimulate the mites hormonal system, entomopathogens are suppressed, and trophobiosis influence on leaves.

Oomen (1982) surveyed predator species and found 22 mite and 2 insect species, listed in Table 8, to prey upon the scarlet mite. From these, the predatory mite *Amblyseius deleoni* is the most common and widespread predator of scarlet mites, although it seems to prefer pink and purple mites to scarlet mites. The population of *A. deleoni* is enhanced by the application of copper fungicides. The predatory mites of the subfamily Stigmaeidae show some preference for the scarlet mite, but also prey on eriophyid and phytoseid mites. They are easier to rear than phytoseid mites. The bug *W. testudo* and the gall midge *Lestodiplosis oomeni* (Table 8) have been identified as the insect predators of the scarlet mite. Nymphs and adults of *W. testudo* are rather selective against the scarlet mite and do not prey on pink or purple mites. They prey on eggs, juveniles and adults of the scarlet mite. The larva of *L. oomeni* preys on purple and pink mites, and also on scarlet mites and its own species when hungry. So far predators have not been mass reared and liberated to control the scarlet mite, but this seems to be a promising biological control method. Control of the scarlet mite without acaricides is more promising with tea grown at higher altitudes (>1000 m) compared to tea at low altitudes.

4 Zusammenfassung

Die Forderung von Tee ohne Rückstände von Pflanzenschutzmitteln steht aus der Sicht des Pflanzenschutzes im Widerspruch zur Forderung des Verbrauchers und des Handels nach hoher Qualität, und der Forderung des Erzeugers nach hohem Ertrag und Verringerung der Arbeitskosten (manuelle Unkrautbeseitigung). Im Teeanbau werden hohe Erträge im Wesentlichen erreicht durch: Beseitigung des Schattens und Verringerung der krankheits- und schädlingsbedingten Verluste. Die Beseitigung der Schattenbäume verändert das AgroÖkosystem mit dem Ergebnis eines erhöhten Unkrautwuchses, des Einsatzes von Düngermitteln und einer Erhöhung der Anfälligkeit gegenüber bestimmten Krankheiten (Sonnenbrand bedingter Stamm-Krebs, Grauer Blattbrand, usw.) und bestimmten Schädlingen (phytophage Milben). Schatten kann jedoch auch ertragsmindernd sein. Bei Auswahl ungeeigneter Schattenbäume oder bei zu hoher Beschattung in der Regenzeit (das Optimum des Lichtgenusses liegt beim Tee zwischen 50% u. 80% des unbehinderten Lichteinfalles). Schatten kann den in Asien auftretenden Befall des Blasenbrandes (*Exobasidium vexans*) und einiger Schädlinge (Tee-Wanze *Helopeltis*) begünstigen. Der wirtschaftliche Schaden durch Krankheiten ist an Tee höher im Vergleich zu Schädlingen, wobei der Blasenbrand u. die Wurzelfäule die wichtigsten Krankheiten sind. Der Befallsdruck von Krankheiten und Schädlingen ist auch von der Bekämpfungsstrategie und den klimatischen Bedingungen abhängig. Letztere ändern sich zwischen den Anbauregionen und mit der Höhenstufe des Teeanbaus. Eine Verringerung des Pflanzenschutzmittelaufwandes kann erreicht werden durch: Verringerung des Befallsdruckes von Krankheiten und Schädlingen, durch Teeanbau unter befallsmindernden Umweltbedingungen (z. B. Höhenstufe, Schattenregulierung), durch Auswahl

krankheits- und schädlingstoleranter Teesorten (Klone), durch die Auswahl von Pflanzenschutzmitteln mit geringen Nebenwirkungen auf natürliche Feinde, und durch Anwendung von Pflanzenschutzmitteln in Abhängigkeit von der wirtschaftlichen Schadschwelle. Spezifische Schadschwellen werden aufgeführt, die meisten müssen noch für die einzelnen Schadorganismen und Anbauggebiete ermittelt werden. Die lange Verweildauer (bis 50 Jahre) des Teebusches an derselben Stelle begünstigt langsam entwickelnden Wurzelkrankheiten, welche zu großen Lichtungen im Teebestand führen und hohe Ernteverluste verursachen. Wurzelkrankheiten werden eingehend besprochen und deren integrierte Bekämpfung mittels Ausmerzungen, dem Einsatz systemischer Fungizide und Bodenfungizide und der Verbesserung der Bodenmikroflora, diskutiert. Ein Überblick der Blatt- und Stammkrankheiten wird gegeben. Die integrierte Krankheits- und Schädlingsbekämpfung des Blasenbrandes und der Scharlachroten Milbe (scarlet mite, *Brevipalpus phoenicis*) werden im einzelnen behandelt.

5 Acknowledgement

The author would like to thank the Tea Research Station Gambung, Java (Indonesia) for invitations to the Station. I am especially indebted to Dr. Attik Dharmadi for the hospitality, repeated intense field visits to tea in Gambung, and the in depth discussion of integrated pest management on tea. Particularly I would like to thank Mrs. Dharmadi for the discussions on root rot of tea.

6 Bibliography and references

- 1 ARUEPRAGASAM, P. V.; ADDAICKAN, S.; KULATUNGA, S. M., 1987: An inexpensive and effective method for the control of red root disease of tea. Sri Lanka Journal of Tea Science 56(1): 5-11
- 2 BARTHAJUR, B. K.; DUTTA, P.; KARAN, SINGH; 1992: Clonal Susceptibility of Red Rust. Two and a Bud 39(2): 52
- 3 BARTHAJUR, B. K.; DUTTA, P.; BEGUM, R.; SARMAH, S. R., 1993: *Bacillus subtilis*, a promising bioagent for control of black rot. Two and a Bud 40(2): 50
- 4 BUSEY, C.F.C.; 1941: The ecology and control of the forest insects of India and the neighbouring countries. Part one: The ecology of forest insects. Forest Research Institute, New Forest, Dehra Dun, India: 767 pp.
- 5 CARR, M. K. V., 1972: The climatic requirements of the tea plant: A review. Experimental Agriculture 8: 1-14
- 6 DAS, S. C., 1974: Parasites and predators on pests of shade trees and ancillary crops in Jorhat circle in north-east India. Two and a Bud 21(1): 17-21 (cited by SARMA, 1979)
- 7 DEKA, M. K.; KARAN SINGH; HANDIQU, R., 1998: Antifeedant and repellent effects of pongam (*Pongamia pinnata*) and wild sage (*Lantana camara*) on tea mosquito bug (*Helopeltis theivora*). Indian J. of Agricultural Sc. 68 (5): 274-276. (cit. Rev. of Agricultural Entomology 86 (10): 9636)
- 8 DHARMADI, A., 1988: Man-initiated factors affecting the population of scarlet mite (*Brevipalpus phoenicis*) in tea plantations in Indonesia. Proceedings of the Symposium on Pest Ecology and Pest Management, Bogor, Indonesia, 9-11 April 1985. SAMEO-BIOTROP special publication No. 32, Bogor, Indonesia: pp. 183-190

- 9 DIHARMAI, A., 1990a: Preying ability of *Mantis hierodula* (Orthoptera: Mantidae) on *Helopeltis antonii* and *Heteropsylla cubana* (in Indonesian language). Pusat Penelitian Gembung (eds): Prosiding Symposium Teh V, Bandung 27 Februar to 1 March: pp. 281-287
- 10 DIHARMAI, A., 1990b: Factors affecting increase of *Helopeltis* in tea plantations (in Indonesian language). Pusat Penelitian Gembung (eds): Prosiding Symposium Teh V, Bandung 27 Februar to 1 March: pp. 175-188
- 11 EDEN, T., 1976: Tea. 3rd edition. Longman, London: 236 pp.
- 12 FAO, 1999: FAO yearbook of production Vol 52, 1998. FAO Statistic Series No. 148. Food and Agriculture Organization of the United Nations, Rome: p. 170
- 13 FUCHS, H. J., 1989: Tea environments and yield in Sri Lanka. Tropical Agriculture 5, Margraf Scientific publisher, Weikersheim (Germany): 320 pp.
- 14 BARTHAJAK, B. K.; DUTTA, P.; BEGUM, R.; SARMAH, S. R., 1993: *Bacillus subtilis*, a promising bioagent for control of black rot. TWD and a Bud 40(2): 50 GADD, C. H., 1949: The commoner diseases of tea. Tea Research Institute of Ceylon, Monographs on Tea Production in Ceylon No. 2 (cit. EDEN, 1967)
- 15 HAINSWORTH, E., 1952: Tea pests and diseases and their control. Heffer, Cambridge
- 16 HAMAYA, E., 1981: Diseases of tea plant in Japan and their control. Rev. Pl. Prot. Res. Japan, 14: 96-111
- 17 HARRIS, K. M., 1982: *Lestodiplosis omenii* sp. n. (Diptera: Cecidomyiidae), a predator on the carinate tea mite *Calcarus carinatus* (Green) (Acarina: Eriophyidae) and other mites on tea plants in Indonesia. Entomologische Berichten 42: 20-24
- 18 KALSHOVEN, L.G.E.; VAN DER LAAN, P.A., 1981: The pests of crops in Indonesia. P.T. Ichtar Baru - Van Hoeve, Jakarta, Indonesia: 701 pp
- 19 KERR, A., SILVA, R.L., 1969: Epidemiology of blister blight (*Exobasidium vexans*). Tea Quart. 40: 1
- 20 KRISHNA, K.; WEESNER, F. M., eds., 1970: Biology of termites. Volume II. Academic Press, New York: 643 pp.
- 21 LEHMANN-DANZINGER, H., 1986: Erste Bestandsaufnahme von Krankheiten an Waldbäumen in Marokko („First evaluation of forest diseases in Morocco,“). Technical report (also available in French). Forest Protection Project PN 81.2005.7-01.200, Deutsche Gesellschaft fuer Technische Zusammenarbeit (GTZ), Eschborn/Ts., Germany: 91 pp.
- 22 LEHMANN-DANZINGER, H., 1987: Spreading mechanism, epidemiology and susceptibility test of Black Sigatoka and Sigatoka diseases. Proceedings Seminar: Improving Citrus and Banana Production through Phytosanitation. 2-5 Dec. 1986, St. Lucia W.I.; CTA-CARDI, Trinidad: pp. 74-94.
- 23 LEHMANN-DANZINGER, H., 1990: Manejo integrado de enfermedades y plagas y diagnóstico de enfermedades en cultivos agrícolas de Costa Rica, 4th ed. (Integrated pest and disease management and diagnostic of diseases of crops in Costa Rica). Technical report CR90-3, PN 80.2054.7.-09.200, Deutsche Gesellschaft fuer Technische Zusammenarbeit (GTZ), Eschborn/TS., Germany: 100 pp.
- 24 LEHMANN-DANZINGER, H., 1994: Introduction to integrated pest management of plant diseases and pests in the tropics and subtropics. 3rd ed. Six Universities Development and Rehabilitation Project (SUDR), Academic Consultant Component (ADB Loan No. 1013-INO), Jakarta, Indonesia. Also published by the Author: Dr. H. Lehmann-Danzinger, Institut f. Pflanzenpathologie u. Pflanzenschutz, Georg-August University of Goettingen, Grisebachstr. 6, 37077 Göttingen, Germany: 182 pp.
- 25 LEHMANN-DANZINGER, H., 1997: Krankheiten an Waldbäumen in den Tropen und Subtropen (Diseases of forest trees in the tropics/subtropics). 3rd ed. Eigenverlag Dr. H. Lehmann-Danzinger, Institut fuer Pflanzenpathologie u. Pflanzenschutz, Georg-August University of Goettingen, 37077 Göttingen, Germany: 219 pp.
- 26 LYR, H., 1995: Modern selective fungicides (properties, applications, mechanisms of action). Gustav Fischer Verlag, Jena - Stuttgart: 595 pp.
- 27 MARSUPONG, M., 1991: Some factors influenced tea resistance to blister blight. Proceedings of the International Symposium on Tea Science, August 26-29, 1991, Shizuoka, Japan: pp.651-654 (cit. Rev. Plant Pathology 73 (1994): 207)

- 28 MULDER, D., 1976: Diseases and disorders of tea (*Camellia sinensis*) in Indonesia. Med. Fac. Landb., Rijksuniv. Gent. 41(2)
- 29 MURALEEDHARAN, V., 1992: Pest control in Asia. Chapter 12, pp. 375-412 in: WILLSON & CLIFFORD
- 30 ONSANDO, J. M.; WARGO, P. M.; WAUDO, S. W., 1997: Distribution, severity, and spread of Armillaria root disease in Kenya tea plantations. Plant Dis. 81: 133-137
- 31 OOMEN, P. A., 1982: Studies on population dynamics of the scarlet mite, *Brevipalpus phoenicis*, a pest of tea in Indonesia. Meddelingen Landbouwhogeschool Wageningen 82-1: 1-88
- 32 PERKOW, W.; PLOSS, H., 1999: Wirksubstanzen der Pflanzenschutz- und Schädlingsbekämpfungsmittel. 3rd ed. Vol. I, Parey Buchverlag, Berlin.
- 33 PITTIARD, A. E., BAKER, E. W., 1958: The false spider mites (Acarina: Tenuipalpidae). University of California Publications in Entomology 14: 175-274
- 34 PURSEGLOVE J. W., 1968: Tropical crops: Dicotyledons 2. Longmans, London: pp. 599-612
- 35 RATTA, P. S., 1992: Pest and disease control in Africa. Chapter 10, pp. 330-352 in: WILLSON & CLIFFORD
- 36 SARMA, P.V., 1979: Possibilities of integrated control of major pests of tea in India. PANS 25: 237-245
- 37 SEMANGUN, H., 1983: Root diseases of tea in Indonesia. J. Plantation Crops (Suppl.), Nov. 1983
- 38 SIANMUGANATHAN, N., 1972: Control of blister blight of tea in Ceylon. South-east Asia (SEA) Regional Symp.Pl.Dis. Tropics, Yogyakarta, Sept. 1972
- 39 SUGHA, S. K., 1997: Perpetuation and seasonal build-up of *Exobasidium vexans*, causal agent of blister blight of tea in Himachal Pradesh. Tropical Science 37: 123-128
- 40 TOMLIN, C. D. S., ed. 1997: The pesticide manual. 11th edition. British Crop Protection Council, Binfield, Bracknell (UK): 1606 pp.
- 41 TUMMAKATE, A.; LEEJIRAJUMNIEN, S., 1980: Tea diseases in Thailand. SEA Symp.Pl. Dis. Tropics II, Bangkok.
- 42 VAN DER GOOT, 1955: Notes on disease incidence of *Exobasidium vexans* on tea. Arch. Theecult. 19: 99-113
- 43 VENKATA RAM, C. S., 1980: Calixin in blister blight control. Agricultural News (BASF) 80(4): 6-7
- 44 WALLER, J. M., HOLDERNESS, M., 1997: Beverage crops and palms. Chapter 9, pp. 229-251 in: Soilborne Diseases of Tropical Crops, HILLOCKS, R. J. & WALLER, J. M., eds., CAB International, Wallingford (UK)
- 45 WILLSON, K. C.; CLIFFORD, M. N., ed. 1992: Tea: cultivation to consumption. Chapman & Hall, London: 769 pp.
- 46 YOUNG, H.E., 1954: Ustilina rot of rubber trees. Rubb. Res. Inst. Ceylon, Adv. Circ. 47